

Are He and N abundances in Type I PNe as high as empirically derived?

Ruth Gruenwald and Sueli M. Viegas

Instituto Astronômico e Geofísico - USP

Av. Miguel Stefano, 4200

04301-904 São Paulo, Brazil

Received _____; accepted _____

ABSTRACT

Type I planetary nebulae (PNe) are defined as those with high He and N abundances (Peimbert & Torres-Peimbert 1983). These objects present in general bipolar geometries and have high stellar temperatures (Corradi & Schwarz 1995, Torres-Peimbert & Peimbert 1997). In this paper we analyse the empirical methods for abundance determination in order to check if the He and N overabundances in Type I PNe are a consequence of a geometrical effect, due to the bipolarity, or the ionization stratification, due to the stellar temperature. For this, we obtain simulated spherically symmetrical as well as bipolar nebulae, using a 3D photoionization code. From the projected emission line intensities for: a) the whole nebula; b) for a slit crossing the nebula; as well as c) for different positions in the nebula, we applied the formulae used in the literature to obtain empirical abundances. These empirical abundances are then compared with the adopted ones. We show that empirical abundances depend on the particular line of sight covered by the observation and can simulate an overabundance and/or the presence of abundance gradients of He and N in planetary nebulae with high stellar temperature. The geometrical effects are also discussed. Systematic errors in abundance determinations by empirical methods are higher for the N/H ratio than for N/O. Thus, it seems better to use the N/O value when discussing N rich objects.

1. Introduction

The knowledge of chemical abundances in Planetary Nebulae (PNe) is important since they can be used as constraints for stellar evolutionary models, as well as for studies of the chemical evolution of galaxies. In fact, PNe abundances reflect the products of stellar evolution, as well as the enrichment of the interstellar medium. Chemical abundances in PNe are usually obtained using empirical methods (as for example in Peimbert & Torres-Peimbert 1987, Kingsburgh & Barlow 1994). Empirical methods are commonly used since they can easily be applied to a great number of objects, from a small number of bright emission lines. Photoionization models also provide the gas chemical composition when applied to specific nebulae (for example, Harrington et al. 1982; Clegg et al. 1987; Gruenwald, Viegas & Broguière 1997).

Following their abundance, PN are classified in types (Peimbert 1978; Peimbert & Torres-Peimbert 1983; Faúndez-Abans & Maciel 1987). Type I PNe are those with high He and N abundances, e.g. $\text{He}/\text{H} \geq 0.125$ and $\log(\text{N}/\text{O}) > -0.30$ (Peimbert & Torres-Peimbert 1983). In a study of morphological and physical properties of PNe, Corradi & Schwarz (1995) state that bipolar nebulae have the hottest stars among PNe, and, except for two objects, all bipolar PNe for which chemical data are available are Type I. Another common characteristic of bipolar nebulae is that the [NII] line intensity is stronger than $\text{H}\alpha$ all over the nebula, apart from the very central region (Corradi & Schwarz 1993b). This characteristic is usually explained by a N overabundance or by shock heating (Corradi & Schwarz 1993a; Corradi & Schwarz 1993c).

Predictions of stellar evolutionary models and the high values obtained for He and N abundances in Type I/bipolar planetaries have suggested that their central stars are more massive than other PN nuclei (Peimbert & Serrano 1980; Calvet & Peimbert 1983; Torres-Peimbert 1984; Corradi & Schwarz 1995). However, new evolutionary models fail to explain

the high abundances of He and N in Type I PNe (Marigo, Bressan & Chiosi 1996), although models including the “hot botton burning” effect and accounting for the chemical evolution on the AGB may lead to higher N/O abundances (van den Hoek & Groenewegen 1997).

From the characteristics of Type I/bipolar planetary nebulae, it can be noted that objects with the highest He and N empirical abundances are those presenting a bipolar geometry and having the highest stellar temperatures. In fact, objects with higher stellar temperatures (T_*) tend to present higher He and N empirical abundances (see Fig. 1). In order to homogenize the data in the figure, only objects with abundances obtained from line intensities integrated in a slit across the nebulae are plotted. As it will be shown in the following sections, empirical abundance determinations depend on the line of sight covered by the observation. Empirical abundances for the objects plotted in Fig. 1 are from de Freitas Pacheco et al. (1992), Kingsburgh & Barlow (1994), and Costa et al. (1996). The stellar temperature corresponds to the Zanstra temperature of HeII. Values for T_* are taken from the literature (Kaler 1983; Preite-Martinez & Pottasch 1983; Pottasch 1984; Shaw & Kaler 1985; Gathier & Pottasch 1988; Gathier & Pottasch 1989; Gleizes, Acker, & Stenholm 1989; Kaler & Jacoby 1989; Shaw & Kaler 1989; Kaler, Shaw, & Kwitter 1990; Stanghellini, Corradi, & Schwarz 1993).

In this paper we analyze the effect of the geometry and of the stellar temperature on the empirical abundance determinations, in order to verify if the high abundances of He and N obtained for Type I PNe are real. In fact, several studies show that the stellar temperature and/or the geometry can affect the abundances obtained from empirical methods. As discussed by Peimbert and collaborators (Peimbert 1967; Peimbert & Costero 1969; Peimbert, Luridiana, & Torres-Peimbert 1995), the gas temperature obtained from observed forbidden line ratios and used in abundance determinations must be corrected by temperature fluctuations inside the nebulae. In fact, for typical conditions of PNe, temperature fluctuations

can be important when the ionizing central star is hotter than 10^5 K (Gruenwald & Viegas 1995). Furthermore, a study on spherically symmetrical homogeneous HII regions show that temperature fluctuations obtained from projected data vary with the line of sight to the nebula; the variation depending on the characteristics of the star and the nebula (Gruenwald & Viegas 1992). With a three-dimensional (3D) self-consistent photoionization code, a model for the bipolar PN IC 4406 has been obtained, showing that its abundance does not fulfill the criteria for Type I, as this nebula is usually classified (Gruenwald et al. 1997). Such an analysis shows that specific and more realistic models which take into account the geometry of the object can lead to abundance results that differ from those obtained with empirical methods.

In order to check for the effects of geometry and stellar temperature on the empirical method to obtain PNe chemical abundances, we apply our self-consistent 3D photoionization code to planetary nebulae. For a wide range of nebular and stellar characteristics, we obtain the physical conditions in each point of the nebula, and calculate the resulting line intensity ratios at different lines of sight to the nebula. From the calculated line intensity ratios we determine the abundances applying the empirical methods used in the literature. These abundances are then compared to those assumed in the models. The models are described in §2. The results are presented in §3 and the conclusions are outlined in §4.

2. Empirical Abundances from Simulated Nebulae

The 3D code has been described in a previous paper (Gruenwald et al. 1997). As in classical uni-dimensional photoionization codes, the input parameters are the stellar temperature and luminosity, as well as the nebular elemental abundances and density distribution. Typical parameters for planetary nebulae are assumed: stellar temperatures (T_*) ranging from 2×10^4 to 3×10^5 K, and stellar luminosities (L_*) from $3 \times 10^2 L_\odot$ to $2 \times 10^4 L_\odot$. The

models are obtained for two sets of chemical abundances (uniform throughout the nebula): solar abundances (Grevesse & Anders 1989), $Z_{\odot}$, and an average abundance for planetary nebulae (Kingsburgh and Barlow 1994), Z_{typ} , which includes He, C, N, O, Ne, S, and Ar; for the remaining elements (Mg, Si, Cl, and Fe), we assume a value equal to a hundredth solar in order to take into account their presence in grains (Stasinska & Tylenda 1986). For spherically symmetrical nebulae, a constant density, n_H , ranging from 10^2 to 10^6 cm^{-3} is assumed. In order to analyze PNe with bipolar appearance, an equatorial torus of denser gas close to the star is included. The presence of a dense equatorial region can explain the observed geometry of bipolar PNe (Calvet & Peimbert 1983; Icke, Preston, & Balick 1989; Sahai et al. 1991; Corradi & Schwarz 1995; Gruenwald et al. 1997). The torus has the following characteristics: i) external radius, r_{ext} , equals to $R/10$, where R is the maximum dimension of the ionized region; ii) internal radius, r_{int} , equal to $r_{ext}/10$; iii) height, h_{disk} , equal to r_{int} ; iv) $n_{torus} = 10 n_H$. The maximum dimension R corresponds to the distance from the star where the fractional ionization of H, H^+/H , equals to 5%, along the torus axis. The torus is centered at the central star. These values for the torus dimensions and density are chosen just to analyze the geometry effect on the abundance determinations, and are not intended to represent all the bipolar nebulae.

The code provides the emissivity of the lines in each point of the nebula. Line intensity ratios can then be obtained: a) for different positions in the projected nebula; b) for the whole nebula; c) for a given slit across the nebula.

We assume that the calculated line intensity ratios simulate observed ratios. Therefore, chemical abundances can be obtained by empirical methods using these line ratios. Comparing the calculated abundances with those adopted for the simulated nebula, the possible effects of geometry and stellar temperature on the empirical abundance calculation can be established. The calculations are performed for different sets of emission line intensity ratios

corresponding to cases (a), (b), and (c), listed above.

In order to obtain empirical elemental abundances, the ionic fractional abundance must be computed. For ions producing detected emission lines, the ion abundance relative to H^+ is obtained from the observed line intensity relative to $H\beta$ and the corresponding line emissivity ratio. Line emissivities depend on the gas density and temperature. The electronic density is obtained from the $[SII] \lambda 6718/\lambda 6733$ line intensity ratio, since this ratio depends weakly on the gas temperature (Osterbrock, 1989). Knowing the density, the gas temperature for the high ionization region ($T_{[OIII]}$) can be obtained from the $[OIII] \lambda\lambda 4959+5007/\lambda 4363$ line ratio, while for the low ionization region ($T_{[NII]}$) from the $[NII] \lambda\lambda 6548+84/\lambda 5755$ line ratio. Line emissivities, electronic density, and temperature are calculated assuming a five-level atom.

A correction must be done for ions which are present in the gas and do not produce detectable emission lines. Such a correction can be done using the corresponding “ionization correction factor” (*icf*) for each element. We apply the *icf* from Peimbert & Torres-Peimbert (1987), as generally used in the literature. Empirical abundances for He, O, and N are then obtained from the following equations:

$$\frac{He}{H} = \frac{He^+ + He^{++}}{H^+} \quad (1)$$

$$\frac{O}{H} = icf(O) \left(\frac{O^+ + O^{++}}{H^+} \right) \quad (2)$$

$$\frac{N}{H} = \left(\frac{N^+}{O^+} \right) \left(\frac{O}{H} \right) = \left(\frac{N^+}{H^+} \right) \left(1 + \frac{O^{++}}{O^+} \right) \left(1 + \frac{He^{++}}{He^+} \right) \quad (3)$$

where

$$icf(O) = \frac{He^+ + He^{++}}{He^+}. \quad (4)$$

The ionization correction factor for O, $icf(O)$, accounts for ionization stages higher than O^{++} . In equation (3), the last two terms correspond to the correction due, respectively, to N^{++} and to higher stages of N.

The abundance of He^+ and He^{++} , relative to H^+ , are determined respectively from the HeI $\lambda 5876$ and HeII $\lambda 4686$ line intensities relative to $H\beta$. The ratio of line emissivities for these lines were calculated following Barker (1978), using the values obtained for the gas density and temperature. A correction due to collisional excitation of He^+ is taken into account following Peimbert & Torres-Peimbert (1987). The O^+ , O^{++} , and N^+ abundances, relative to H^+ , are obtained respectively from $[OII] \lambda\lambda 3727+29/H\beta$, $[OIII] \lambda\lambda 4959+5007/H\beta$, and $[NII] \lambda\lambda 6548+84/H\beta$ line ratios.

3. Results

In the following, the emission line ratios from simulated nebulae and the corresponding empirical chemical abundances are discussed for cases (a), (b), and (c) listed above.

3.1. Spherically symmetrical nebulae

The behavior of the $[NII] \lambda\lambda 6548+84/H\alpha$ line intensity ratio with the projected distance to the center of the nebula is shown in Fig. 2a, while Fig. 2b gives the corresponding N/H abundance ratio. The simulations are performed with uniformly distributed density equal to 10^2 cm^{-3} , $Z = Z_{typ}$, and stellar luminosity $L_* = 3 \times 10^3 L_\odot$. The projected distance to the center of the nebula is given in units of the radius of the ionized region. The curves are labeled by the stellar temperature, in units of 10^3 K . The $[NII]/H\alpha$ line ratio depends on the position of the line of sight to the nebula since different lines of sight cross different proportions of $[NII]$ and $H\alpha$ emitting regions (Fig. 2a). The behavior of the $[NII]/H\alpha$ line ratio for different

lines of sight depends on the stellar temperature. Higher stellar temperatures lead to larger recombination regions, increasing the H^+ emitting volume relative to N^+ emitting volume. Therefore, there is a shift of the peak of $[NII]/H\alpha$ towards smaller r_{proj}/R with increasing T_* , as shown in Fig. 2a. The maximum value for $[NII]/H\alpha$ depends on the value of the nitrogen abundance used in the models. However, as seen in the figure, the measured line ratio can substantially vary even for a nebula with a homogeneous abundance.

Details on the $[NII]/H\alpha$ distribution on the projected nebula depend also on the stellar luminosity, on the nebular density, as well as on the chemical abundance. For increasing values of the nebular density, the region where $[NII]/H\alpha$ is higher than one are relatively narrower with the peak shifted to higher r_{proj} . The peak is also shifted to higher r_{proj} for increasing L_* . Lower values for the iron abundance increases the $[NII]/H\alpha$ ratio since Fe is an important cooler in the region where $[NII]$ lines are produced. This effect is more significant for lines of sight far from the central regions.

Following Corradi & Schwarz (1993b), $[NII]$ stronger than $H\alpha$ appears to be a common characteristic of bipolar nebulae. Our results show that even for a nebula with solar N/H abundance (which is lower than the average N/H for PNe), the measured $[NII]/H\alpha$ line ratio can be higher than one. Indeed, a simulated nebulae with solar abundances and high T_* (typical of Type I planetary nebulae) present $[NII]/H\alpha$ higher than one all over the nebula. For NGC 2440, which is a PN with a high value for T_* , the $[NII]$ line is brighter than $H\alpha$, and the line ratio increases from 3 to 30, from the central to the outer zones (Icke et al. 1989). In fact, using characteristic parameters for this object ($T_* = 2 \times 10^5$ K, $L_* = 5 \times 10^3 L_\odot$, and $n_H = 6 \times 10^3 \text{ cm}^{-3}$), and for $Z = Z_{typ}$, we verify that the $[NII]/H\alpha$ ratio can reach values of the order of 26. Notice that these values are given just as an illustration, since these results correspond to a spherical symmetrical nebula, while NGC 2440 is clearly irregular (see, for example, Schwarz, Corradi & Melnick 1992).

The variation of the empirical N abundance in different lines of sight to a simulated nebula, is shown in Fig. 2b. The empirical values of N/H are plotted relative to the adopted abundance ratio. The empirical value obtained for the abundance depends on the detailed ionization structure of the nebula. Notice that for PNe whose central stellar temperature is high ($T_* > 10^5$ K), the obtained values for different lines of sight can significantly vary, and can be higher than the adopted one in a extended region, mimicking an unreal overabundance, as well as an abundance gradient. Therefore, studies of abundance gradients of nitrogen across a nebula applying empirical methods to the observed line intensities (Chu et al. 1991; Guerrero, Stanghellini, & Manchado 1995; Guerrero, Manchado, & Serra-Ricart 1996; Guerrero 1997; Corradi et al. 1997) can be misleading. For $T_* \leq 10^5$ K, the resulting abundances are not very different from the adopted one, except in a narrow region near the borders. For such stellar temperatures, the volume of the H^o zone is larger than the N^o one. Lines of sight with increasing distance from the center include a higher fraction of low ionized regions. Therefore, empirical methods which do not account for neither N^o nor H^o overestimates the N abundance and must not be applied in this case. For models with different n_H and/or L_* , the discrepancies from the adopted abundance are somewhat different. However, the major effect comes from the stellar temperature.

The variation of the $[NII]/H\alpha$ with T_* is shown in Fig. 3a for the whole nebula and for a long and narrow slit across its central part. The two upper curves correspond to simulated nebulae with Z_{typ} , while the lower ones to $Z_\odot$. The long-dashed and solid lines represent line ratios for the entire nebula, while the dot-dashed and short-dashed curves correspond to line intensities integrated in a slit. The nebular density and the stellar luminosity are the same as in Fig. 2. As expected, $[NII]/H\alpha$ values are lower for a slit than for the whole nebula. In fact, in the latter case there is a higher contribution of low ionized regions, increasing the $[NII]$ contribution relative to $H\alpha$, for a given T_* .

Results for the empirical N/H relative to the adopted abundance are shown in Fig. 3b, for the case $Z = Z_{\odot}$. For $Z = Z_{typ}$, the results are similar, except for $T \leq 40000$ K because of the effect of Fe on the gas cooling. The curves in Fig. 3b show that the empirical N/H abundances are overestimated for $T_* > 8 \times 10^4$, while for lower T_* the N abundance is lower than the true value. This behavior is related to the ionization correction factors used in the empirical calculations (see equations [1] to [4]). In fact, for $T_* < 8 \times 10^4$ K the contribution of N^{++} is underestimated since the correction for this non-observed ion is based on O^{++} which has a smaller emitting volume than N^{++} (the volume of the O^+ zone being greater than the N^+ one due to charge transfer reactions). The main effect for high T_* is that the He^{++} zone contains the N^{++} zone; so, when applying the ionization correction factor for N^{++} , this ion is taken into account twice (both from the O^{++} and He^{++} corrections), increasing the obtained empirical N/H. Results for N/H from emission line intensities obtained with a slit crossing the nebular central region are greater than those where the whole nebula is observed. This comes from the fact that with such a slit the observed emitting volume of high ionized zones relative to low ionized zones is larger than for the whole nebula.

The behavior of the O/H empirical abundances with T_* is similar to that of N/H, although less affected by the geometry. For example, when compared to N, the O abundance becomes greater than the true value for higher stellar temperatures. This leads to discrepancies of N/O which increase with T_* (Fig. 3b).

Since He/H is one of the criteria for classifying Type I PNe, this ratio is shown in Fig. 4. For He/H, empirical methods do not take into account H^o nor He^o abundances. However, for $T_* < 4 \times 10^4$ K, there is an increasing He^o zone, larger than the H^o one; when He^o is not taken into account, lower abundances for He are then obtained. For $T_* > 10^5$ K, the H^o zone is increasingly larger than the He^o one due to the high energy photons which can ionize He^o very far into the nebula. If this effect is not properly taken into account, larger He/H are

obtained.

For increasing values of n_H , the discrepancies of empirical values of N/H increase, while those of He/H decrease (for increasing densities the volume of He^o and H^o tend to be equal). For example, for $n_H = 10^4 \text{ cm}^{-3}$, and same L_* as in Fig. 3b, the empirical N/H ratio is higher by a factor more than 3 for $T_* \geq 2 \times 10^5 \text{ K}$. Regarding the stellar luminosity, the effect is practically negligible for N, but the discrepancies of He/H increase for lower L_* .

3.2. Bipolar nebulae

The effect of the torus and of the resulting bipolarity on the $[\text{NII}]/\text{H}\alpha$ line ratio and on the corresponding empirical N/H are shown respectively in Fig. 5a and 5b. The curves correspond to $T_* = 10^5 \text{ K}$, $L_* = 3 \times 10^3 L_\odot$, and $Z = Z_{typ}$. Solid lines correspond to a spherically symmetrical nebula with a uniform density ($n_H = 10^2 \text{ cm}^{-3}$) throughout the nebula. Dashed and dot-dashed curves show the result for a nebula with the same chemical abundance, T_* , L_* , and overall density, but including a torus around the central star. The torus density is $n_{torus} = 10n_H$, and its dimensions are as described in §2. The dashed curve corresponds to a long slit along the lobes, and the dot-dashed one to a slit perpendicular to the torus axis, both passing through the central star. For a bipolar nebula the $[\text{NII}]/\text{H}\alpha$ ratio as well as the resulting N/H vary with the position of the line of sight to the nebula, due to its geometry. However, the effect due to the ionization distribution inside the nebula determined by the stellar temperature is much stronger, when the observations are made along the lobes, as it is often the case. For a given set of T_* , L_* , and overall density n_H , the main differences between the results for a bipolar and a homogeneous nebula are related to the torus characteristics, i.e, the contrast of n_H with the torus density, as well as to the torus dimensions. For a slit along the maximum extent of the lobes, the main differences appear for lines of sight crossing the torus. The opening angle of the torus will be important

for slits which make an angle with the torus axis. Any intermediate slit position will show a variation between the dashed and dotted-dashed curves of Fig. 5a and 5b. The results presented in these figures are an illustration of the ambiguity on the determination of any abundance gradients using empirical methods on bipolar objects.

4. Conclusions

The results of the preceding section show that empirical methods for abundance determination can simulate an overabundance and/or the presence of abundance gradients of He and N in planetary nebulae with high T_* , even when applied to spherically symmetrical and homogeneous objects. This is also true for N/O. Recall that Type I PNe, defined as those nebulae which present an overabundance of He and N, have the hottest stars among PNe, and are generally bipolar. As shown in §3, both characteristics can induce an apparent overabundance when empirical methods are used.

Following our results (Figs. 3a and 3b), systematic errors in the abundance determinations by empirical methods are smaller for N/O. Thus, it is better to use the N/O ratio instead of N/H when discussing N rich planetaries. For central stellar temperatures in the range 10^5 to 3×10^5 K the N/O ratio can be overestimated by a factor of 1.5. On the other hand, the average N/O value for Type I PNe is about 2.8 higher than Type II PNe (Peimbert 1990). So, we expect that, even with a better N/O determination, Type I PNe may still be N rich compared to Type II PNe.

Inference about abundances on planetary nebulae are important tests to stellar evolutionary models. Several arguments indicate that Type I PNe come from more massive progenitors (Torres-Peimbert & Peimbert 1997). In particular, high masses for the central stars of Type I PNe are generally invoked to explain their high He and N abundances since

the higher N/O ratio, the higher the original mass of the central star. Some theoretical evolutionary models fail to explain these overabundances (Marigo et al. 1996). However, models accounting for the chemical evolution on AGB and including the "hot bottom burning" effect are consistent with PNe abundances (van den Hoek & Groenewegen 1997). Our results show that He and N overabundances in Type I PNe may be lower than indicated by the empirical methods. The correction of the empirical abundances by those systematic errors change the constraints to the evolutionary models and may lead to different conclusions on the importance of the different processes.

Furthermore, from the discussion in the preceding section, empirical methods applied to PNe with high stellar temperatures can mimic abundance gradients. Thus abundance gradients and their relation to multiple ejection phenomena must also be reviewed.

Let us recall that observed line intensities are integrated on the line of sight, and different lines of sight cross different fractions of regions differing by their ionization state. Since lines originate in different regions, the observed image of a nebula depends on choice of the emission line. As shown in the preceding section, abundances obtained from observed line intensities depend on the line of sight. Therefore, empirical abundances obtained from line intensities averaged over different points in a nebula, or conclusions based on average abundances for a given nebula, may be incorrect.

The effects shown in the present paper must be taken as a warning when discussing chemical enrichment and/or abundance gradients in a given nebula. Any conclusion about abundances, in particular in nebulae with a high star temperature, even with simple geometries, must rely on detailed photoionization models which properly account for the integration of emission line intensities on a given line of sight. Classical unidimensional photoionization models are suitable for spherically symmetrical objects, whereas a 3D photoionization code (Gruenwald et al. 1997) must be used for other geometries.

We are thankful to M. Peimbert for stimulating discussions and valuable suggestions. This work was partially supported by Fapesp, CNPq, and PRONEX/FINEP.

REFERENCES

- Barker, T. 1978, *ApJ*, 220, 193
- Calvet, N., & Peimbert, M. 1983, *Rev. Mex. Astron. Astrof.*, 5, 319
- Chu, Y.-H., Manchado, A., Jacoby, G. H., & Kwitter, K. B. 1991, *ApJ*, 376, 150
- Clegg, R. E. S., Harrington, J. P., Barlow, M. J., & Walsh, J. R. 1987, *ApJ*, 314, 551
- Corradi, R. L. M., & Schwarz, H. E. 1993a, *A&A*, 268, 714
- Corradi, R. L. M., & Schwarz, H. E. 1993b, *A&A*, 269, 462
- Corradi, R. L. M., & Schwarz, H. E. 1993c, *A&A*, 278, 247
- Corradi, R. L. M., & Schwarz, H. E. 1995, *A&A*, 293, 871
- Corradi, R. L. M., Perinotto, M., Schwarz, H. E., & Claeskens, J. F. 1997, *A&A*, 322, 975
- Costa, R. D. D., Chiappini, C., Maciel, W. J., & de Freitas Pacheco, J. A. 1996, *A&A*, 116, 249
- de Freitas Pacheco, J. A., Maciel, W. J., & Costa, R. D. D. 1992, *A&A*, 261, 579
- Faúndez-Abans, M., & Maciel, W. J. 1987, *A&A*, 183, 324
- Gathier, R., & Pottasch, S. R. 1988, *A&A*, 197, 266
- Gathier, R., & Pottasch, S. R. 1989, *A&A*, 209, 369
- Gleizes, R. S., Acker, A., & Stenholm, B. 1989, *A&A*, 222, 237
- Grevesse, N., & Anders, E. 1989, in *Cosmic Abundances of Matter*, Proceedings of the AIP Conference (American Institute of Physics), 1
- Gruenwald, R., & Viegas, S. M. 1992, *ApJS*, 78, 153
- Gruenwald, R., & Viegas, S. M. 1995, *A&A*, 303, 535
- Gruenwald, R., Viegas, S. M., & Broguière, D. 1997, *ApJ*, 480, 283

- Guerrero, M. A. 1997, thesis, Instituto de Astrofisica de Canarias
- Guerrero, M. A., Manchado, A., & Serra-Ricart, M. 1996, *ApJ*, 456, 651
- Guerrero, M. A., Stanghellini, L., & Manchado, A. 1995, *ApJ*, 444, L49
- Harrington, J. P., Seaton, M. J., Adams, S., & Lutz, J. H. 1982, *MNRAS*, 199, 517
- Icke, V., Preston, H. L., & Balick, B. 1989, *AJ*, 97, 462
- Kaler, J. B. 1983, *ApJ*, 271, 188
- Kaler, J. B., & Jacoby, G. H. 1989, *ApJ*, 345, 871
- Kaler, J. B., Shaw, R. A., & Kwitter, K. B. 1990, *ApJ*, 359, 392
- Kingsburgh, R. L., & Barlow, M. J. 1994, *MNRAS*, 271, 257
- Marigo, P., Bressan, A., & Chiosi, C. 1996, *A&A*, 313, 545
- Osterbrock, D. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (University Science Books)
- Peimbert, M. 1967, *ApJ*, 150, 825
- Peimbert, M. 1978, *Planetary Nebulae, Proceedings of the Symposium*, Ithaca, Dordrecht, D. Reidel Publishing, Co., 215
- Peimbert, M. 1990, *Rep. Prog. Phys.*, 53, 1559
- Peimbert, M., & Costero 1969, *Bol. Obs. Ton. y Tac.*, 5, 3
- Peimbert, M., Luridiana, V., & Torres-Peimbert, S. 1995, *Rev. Mex. Astron. Astrof.*, 31, 147
- Peimbert, M., & Serrano, A. 1980, *Rev. Mex. Astron. Astrof.*, 5, 9
- Peimbert, M., & Torres-Peimbert, S. 1983, in *IAU Symp. 103, Planetary Nebulae*, ed. D. R. Flower (Dordrecht: Kluwer), 233
- Peimbert, M., & Torres-Peimbert, S. 1987, *Rev. Mex. Astron. Astrof.*, 14, 540

- Pottasch, S. R. 1984, *Planetary Nebulae* (Dordrecht: Reidel)
- Preite-Martinez, A., & Pottasch, S. R. 1983, *A&A*, 126, 31
- Sahai, R., Wooten, A., Schwarz, H. E., & Clegg, R. E. S. 1991, *A&A*, 251, 560
- Schwarz, H. E., Corradi, R. L. M., & Melnick, J. 1992, *A&AS*, 96, 23
- Shaw, R. A., & Kaler, J. B. 1985, *ApJ*, 295, 537
- Shaw, R. A., & Kaler, J. B. 1989, *ApJS*, 69, 495
- Stanghellini, L., Corradi, R. L. M., & Schwarz, H. 1993, *A&A*, 276, 463
- Stasinska, G., & Tylenda, R. 1986, *A&A*, 155, 137
- Torres-Peimbert, S. 1984, *Stellar Nucleosynthesis*, eds. C. Chiosi and R. Renzini, D. Reidel Publishing Co., 3
- Torres-Peimbert, S., & Peimbert, M. 1997, in *IAU Symp. 180, Planetary Nebulae*, in press
- van den Hoek, L. B., & Groenewegen, M. A. T. 1997, *A&AS*, 123, 305

Fig. 1.— Empirical abundance ratios from the literature (see text) versus the stellar temperature for N/H (a) and He/H (b).

Fig. 2.— [NII]/H α line ratio (a) and the empirical N/H (b) relative to the adopted value as a function of the line of sight. The projected distance to the center of the nebula is given in units of the nebula ionized radius. Curves correspond to simulated spherically symmetrical nebulae with $L_* = 3000 L_\odot$, $n_H = 10^2 \text{ cm}^{-3}$, and Z_{typ} , and are labeled by the stellar temperature in units of 10^3 K .

Fig. 3.— (a) [NII]/H α versus T_* . The two upper curves correspond to Z_{typ} , while the lower ones to $Z_\odot$. Long-dashed and solid lines represent line ratios for the whole nebula, while dot-dashed and short-dashed curves correspond to line intensities integrated in a slit. The values of L_* and n_H are the same as in Fig. 2. (b) empirical N/H and N/O relative to the adopted value ($Z_\odot$).

Fig. 4.— same as Fig. 3b, for He/H

Fig. 5.— [NII]/H α (a) and N/H (b) relative to the adopted ratio as a function of the line of sight, for $T_* = 10^5 \text{ K}$. Solid lines correspond to a spherically symmetrical nebula, while dashed and dot-dashed correspond to a bipolar nebula of same stellar temperature, respectively for a slit along the lobes, and a slit perpendicular to the torus axis.









